Application No: 09/831,272 Response to Office Action mailed May 28, 2008 Docket No. 9730-1

Page 2 of 10

AMENDMENTS TO THE CLAIMS

This complete listing of claims will replace all prior versions, and listings, of claims in the application:

We claim:

- 1. (Canceled)
- (Currently Amended) A chimeric promoter capable of local gene expression in
 plants wherein the expression is induced by elicitor treatment, pathogen infection, or both
 comprising:
 - (i) two or more cis-acting elements sufficient to direct: pathogen-elicitor-specific expression of a nucleic acid sequence, pathogen-infection-specific expression of a nucleic acid sequence, or both, wherein at least one of the said two or more cis-acting elements consists of the nucleotide sequence of SEQ ID NO: 11, and
 - (ii) a minimal promoter,

wherein induction of said local gene expression upon elicitor treatment or pathogen infection is between 10-fold and 15-fold.

 (Currently Amended) A chimeric promoter capable of local gene expression in plants wherein the expression is induced by elicitor treatment, pathogen infection, or both, Application No: 09/831,272 Response to Office Action mailed May 28, 2008 Docket No. 9730-1

Page 3 of 10

comprising:

- (i) two or more cis-acting elements sufficient to direct: pathogen-elicitor-specific expression of a nucleic acid sequence, pathogen-infection-specific expression of a nucleic acid sequence, or both, wherein at least one of the said two or more cis-acting elements consists of the nucleotide sequence of SEQ ID NO: 11 and
 - (ii) a minimal promoter,

further comprising a *cis*-acting element having the nucleotide sequence selected from the group consisting of: SEO ID NO: 1 and SEO ID NO: 2.

- 4-7. (Canceled)
- 8. (Currently Amended) The chimeric promoter of claim 2, 3, 39, 42, 43, 47 or 49, wherein a spacer region composed of 4 to 10 base pairs separates at least two of said cisacting elements are separated by a spacer of between about 4 to 10 base pairs.
- 9. (Currently Amended) The chimeric promoter of claim 2, 3, 39, 42, 43, 47 or 49, wherein at least two of said two or more *cis*-acting elements are separated by a spacer of between about 50 to 1000 base pairs.
 - 10-21. (Canceled)

{WP537716;1}

Application No: 09/831,272 Response to Office Action mailed May 28, 2008 Docket No. 9730-1

Page 4 of 10

 (Previously Presented) An isolated cis-acting element sufficient to direct pathogen-elicitor-specific expression, pathogen-infection-specific expression, or both, consisting of the nucleotide sequence of SEQ ID NO: 11.

23-38. (Canceled)

39. (Currently Amended) The A promoter obtainable by a method of rendering a gene responsive to pathogens comprising inserting at least one *cis*-acting element sufficient to direct pathogen-elicitor-specific expression, pathogen-infection-specific expression, or both, into the promoter of said gene, wherein (1) induction of local gene expression in plants upon elicitor treatment, pathogen infection, or both, is between 10-fold and 15-fold and wherein the at least one *cis*-element promoter comprises SEQ ID NO: 11, or (2) wherein the induction of local gene expression in plants is at least 15-fold and the at least one *cis*-acting element promoter comprises two copies of SEQ ID NO: 11, or a combination of one copy of SEQ ID NO: 11 and one copy of SEO ID NO: 7.

40-41. (Canceled)

42. (Currently Amended) A chimeric promoter capable of local gene expression in plants wherein the expression is induced by elicitor treatment, pathogen infection, or both, comprising:

(WP537716:1)

Page 5 of 10

- (i) two or more cis-acting elements sufficient to direct: pathogen-elicitorspecific expression of a nucleic acid sequence, pathogen-infection-specific
 expression of a nucleic acid, or both, wherein at least one of the two or more cisacting elements eonsists comprise at least one of the nucleotide sequence of SEQ ID
 NO: 11; and wherein at least one of the two or more cisacting elements consists of
 the nucleotide sequence of SEO ID NO: 7, and
- (ii) a minimal promoter.
- 43. (Previously Presented) The chimeric promoter according to claim 42, wherein the two or more cis-acting elements comprise two copies of the nucleotide sequence of SEQ ID NO: 11 and two copies of the nucleotide sequence of SEO ID NO: 7.
- (Currently Amended) A recombinant gene comprising the chimeric promoter of claim
 3, 8, 9, 39, 42, or elaim 43, 47 or 49.
- (Currently Amended) A vector comprising the chimeric promoter of claim 2, 3, 8,
 39, 39, 42, er-elaim 43, 47 or 49.
- 46. (Withdrawn Currently Amended) A method for the production of transgenic plant transgenic plant tissue comprising the introduction of introducing



a chimeric promoter according to claim 2, 3, 8, 9, 39, 42, er 43, 47 or 49, into the genome of said plant, plant cell or plant tissue.

- 47. (Currently Amended) A chimeric promoter capable of local gene expression in plants, wherein the expression is induced by elicitor treatment, pathogen infection, or both, comprising:
- two or more cis-acting elements sufficient to direct: pathogen-elicitorspecific expression of a nucleic acid sequence, pathogen-infection-specific expression of a nucleic acid sequence, or both, and
 - (ii) a minimal promoter,

wherein induction of said local gene expression upon elicitor treatment or pathogen infection is at least 15-fold, the chimeric promoter comprising:

two copies of SEQ ID NO: 11, or

the combination of one copy of SEQ ID NO: 11 followed by one copy of SEQ ID NO: 7, or

the combination of four copies of SEQ ID NO: 11 followed by four copies of SEO ID NO: 7.

- 48. (Canceled)
- 49. (Currently Amended) A chimeric promoter capable of local gene expression in

Application No: 09/831,272 Response to Office Action mailed May 28, 2008 Docket No. 9730-1

Page 7 of 10

plants, wherein the expression is induced by elicitor treatment, upon pathogen infection comprising:

- (i) two or more cis-acting elements sufficient to direct: pathogen-elicitor-specific expression of a nucleic acid sequence, pathogen-infection-specific expression of a nucleic acid, or both, wherein at least one of the two or more cisacting elements consists of the nucleotide sequence of SEQ ID NO: 11 and
- (ii) a minimal promoter,

further comprising a cis-acting element having the nucleotide sequence selected from the group consisting of: SEQ ID NO: 3 and SEQ ID NO: 4.